

Vacuum Stability of a General Scalar Potential of a Few Fields

Kristjan Kannike

National Institute of Chemical Physics and Biophysics, R  vala 10, 10143 Tallinn, Estonia

kannike@cern.ch

Abstract

We calculate analytical vacuum stability or bounded from below conditions for general scalar potentials of a few fields. After a brief review of copositivity, we show how to find positivity conditions for more complicated potentials. We discuss the vacuum stability conditions of the general potential of two real scalars, without and with the Higgs boson included in the potential. As further examples, we give explicit vacuum stability conditions for the two Higgs doublet model with no explicit CP breaking, and for the \mathbb{Z}_3 scalar dark matter with an inert doublet and a complex singlet. We give a short overview of positivity conditions for tensors of quartic couplings via tensor eigenvalues. A Mathematica notebook is included with the source files.

Contents

1	Introduction	2
2	Copositivity & Orbit Spaces	4
2.1	Copositivity	4
2.2	Orbit Spaces	5
3	Vacuum Stability of the Scalar Potential of Two Real Scalars	6
3.1	Vacuum Stability Conditions from Positivity of a Quartic Polynomial	6
3.2	Bounded From Below Conditions for a Quartic Polynomial on \mathbb{R}_+	8
3.3	Vacuum Stability Conditions from Positivity with an Affine Space	9
4	Vacuum Stability for Two Real Scalars & the Higgs boson	10
5	Vacuum Stability of the 2HDM with Real Couplings	14
6	Vacuum Stability for \mathbb{Z}_3 Scalar Dark Matter	17
7	Tensor Eigenvalues	19
7.1	Positive Tensors	19
7.2	Vacuum Stability of the Potential of Two Real Scalars	20
8	Conclusions	21

1 Introduction

A scalar potential has to be bounded from below to make physical sense. In the Standard Model (SM), it simply means that the self-coupling of the Higgs boson has to be positive. In an extended model with more scalar fields, the potential has to be bounded from below – the vacuum has to be stable – in the limit of large field values in all possible directions of the field space. In this limit, any terms with dimensionful couplings – mass or cubic terms – can be neglected in comparison with the quartic part of the scalar potential.¹

In quantum field theories, scalar couplings change with energy due to the renormalisation group running. The vacuum stability conditions may be satisfied at some scales and not satisfied at others. Checking the vacuum stability of the tree level potential with running couplings can help to determine the scale of validity of a given model. On the other hand, in models with classical scale invariance [1], where tree-level mass terms are absent, violation of vacuum stability conditions at a *finite* field range can be used to produce minima and induce symmetry breaking via dimensional transmutation as in the Coleman-Weinberg mechanism [2].

The remarkable Tarski-Seidenberg theorem implies that the question of whether the vacuum is stable or not for given values of the scalar couplings is in principle always decidable. Nevertheless, the general problem of finding whether a given polynomial is non-negative is an NP-hard problem if the degree of the polynomial is at least four [3], which is the case for renormalisable scalar potentials in four dimensions.

The most general quartic potential of real scalars is, of course,

$$V(\phi) = \lambda_{ijkl} \phi_i \phi_j \phi_k \phi_l, \quad (1)$$

where the coupling tensor λ_{ijkl} can be always made completely symmetric under any exchange of the indices. Alas, relatively simple complete conditions of the positivity of the potential (1) can be given only in the case of two fields (Sect. 3), or of three fields, if the potential is biquadratic in one of them (Sect. 4).

Since $V(\phi)$ is a homogenous quartic polynomial, scaling the fields by a positive real constant c gives $V(c\phi) = c^4 V(\phi)$ and does not affect vacuum stability.² Therefore, we can write the quartic potential as

$$V(\phi) = V(\hat{\phi}) r^4, \quad (2)$$

where $\hat{\phi}^2 = 1$ and $r \geq 0$. We see that if $V(\hat{\phi})$ is negative for some $\hat{\phi}$, then the potential tends to negative infinity as $r \rightarrow \infty$ and the vacuum is not stable.

Thus, to determine whether a potential is bounded from below in the limit of large field values, we can minimise its quartic part on a unit hypersphere, enforced by a Lagrangian multiplier λ :

$$V(\phi, \lambda) = V(\phi) + \frac{\lambda}{2} (1 - \phi^2), \quad (3)$$

which yields the stationary point equations

$$\frac{\partial V(\phi)}{\partial \phi_i} = \lambda \phi_i, \quad \phi^2 = 1. \quad (4)$$

¹The requirement of strong stability means demanding that the quartic part of the potential $V_4 > 0$ as the fields $\varphi_i \rightarrow \infty$, whereas $V_4 \geq 0$ gives stability in the marginal sense and there can be flat directions (then the quadratic or mass squared part of the potential *has* to be positive and there cannot be any cubic terms). For simplicity, we give conditions for strong stability, which in practice means making inequalities strict.

²We can even scale each field ϕ_i separately by a different positive coefficient. In particular, we can scale it by $\phi_i \rightarrow \phi_i / \lambda_i^{\frac{1}{4}}$, where λ_i is its self-coupling, and make the coefficient of ϕ_i^4 equal to unity for the purpose of calculating positivity conditions (this only holds at tree level or at a fixed energy scale).

Notice that $\lambda = 4V_{\min}$ since if we write the constraint on ϕ as $g(\phi, c) = c^2 - \phi^2$, then on one hand $dV_{\min}/dc = \lambda c$, while on the other hand $dV_{\min}/dc = 4c^3V_{\min}$ with $c = 1$ in the end.

Global or gauge symmetries of the potential may help to simplify the problem. An important special case is given by quartic potentials that are biquadratic in fields and have the form

$$V = \lambda_{ij} \phi_i^2 \phi_j^2. \quad (5)$$

The couplings λ_{ij} can be written as a matrix on the basis of ϕ_i^2 . Since the squares of real fields are non-negative, the natural domain of such potentials is not \mathbb{R}^n but the non-negative orthant \mathbb{R}_+^n . Positivity on \mathbb{R}_+^n or copositivity (short for ‘conditional positivity’) was introduced in [4].³ Copositivity has found wide use in the field of convex optimisation, and was first used to derive vacuum stability conditions in [6]. The set of copositive matrices is larger than and includes the familiar set of positive semidefinite matrices.⁴ While the positive definite part of the parameter space can be easily found via Sylvester’s criterion [8], the criteria for copositivity are more involved, but definite analytic procedures exist to compute them.

Even if the fields are gauge multiplets, any potential can be written in terms of field magnitudes and orbit space variables [9, 10, 11, 12, 13]. In many cases the potential is a monotonous function of orbit space parameters and its minimum occurs on the boundary of the orbit space. The symmetries of the potential may restrict the variables to a more complicated space such as the ‘future light cone’ orbit space of the two Higgs doublet model (2HDM) [14, 15, 16, 17] or the similar orbit spaces of the 3HDM [18] and NHDM [19]. The importance of taking the orbit space into account properly can be seen, for example, in the case of the type II seesaw: the vacuum stability conditions calculated in [20] were somewhat too strong, because the orbit space parameters do not vary independently [21].

In general, the conditions for a potential to be bounded from below can be expressed in many ways. It may be possible to produce conditions for vacuum stability that are analytical but of considerable length. For more complicated potentials with several fields, one has to resort to numerics, for which the methods we present can still be useful for reducing the parameter space to scan over. The purpose of the paper is to introduce into the ‘toolbox’ of calculating vacuum stability conditions some methods that are specific, but useable in many practical cases, and others that are more complicated but also more general. While recent mathematical literature is concerned with approximate methods of finding positivity for polynomials of many variables, particle physics models usually deal with a few scalar fields and analytical solutions may afford more insight.

On numerous occasions, the new addition to the scalar sector consists of just a couple of real scalar singlets, often in the guise of a complex singlet. In this case the vacuum stability reduces to the problem of positivity of a general quartic polynomial. For comparison, we also derive the vacuum stability conditions in another form, using the Sylvester criteria for the matrix of scalar couplings. And, of course, no low energy scalar potential is complete without the Standard Model Higgs doublet, which we learn to include as well. Similar conditions can be derived e.g. for the 2HDM, where the potential can be considered to be a quartic polynomial in magnitudes of fields, or for more complicated models, such as the \mathbb{Z}_3 scalar dark matter [22, 23] with an inert doublet and a complex singlet. To our knowledge, the results for the potentials of the two singlets (and the Higgs) and for the \mathbb{Z}_3 scalar dark matter are new. As for the 2HDM with real couplings, our results are in shorter form than similar results in the literature [24].

In addition, we reconsider the notion of copositivity of matrices and discuss its relation to orbit space variables. In more complicated situations, tests of (co)positivity in terms of eigenvalues of the *tensor* of scalar couplings can help. Similarly to a positive matrix, a tensor is positive-definite if its eigenvalues associated with its real eigenvectors are positive.

In Sect. 2 we give a brief review of orbit spaces and copositivity. In Sect. 3 we give the conditions for the general potential of two real scalars to be positive. It is not too hard to include the SM Higgs

³See [5] for a good review of copositive matrices.

⁴For a recent review on positive semidefinite matrices, see [7].

doublet into the potential in Sect. 4. In Sect. 5 we derive vacuum stability conditions for the 2HDM with no explicit CP-breaking. Sect. 6 provides another illustration in the vacuum stability conditions for \mathbb{Z}_3 scalar dark matter. In Sect. 7 we introduce tensor eigenvalues as a way to determine the vacuum stability conditions for a most general scalar potential. We conclude in Sect. 8. A Mathematica notebook with the conditions for all examples is included with the L^AT_EX source of the paper.

2 Copositivity & Orbit Spaces

A scalar potential (5) biquadratic in fields is bounded from below if the matrix of couplings λ_{ij} is copositive [6]. Even if the fields are higher multiplets under a gauge group, any potential can be written in terms of squares of field magnitudes and a few dimensionless orbit space variables.

2.1 Copositivity

The criteria to determine whether a matrix is positive in the usual sense are well established. *A symmetric matrix A is said to be positive semidefinite if the quadratic form $x^T A x \geq 0$ for all vectors x in \mathbb{R}^n . A symmetric matrix A is positive definite if the inequality is strict, $x^T A x > 0$ for any non-zero vector x in \mathbb{R}^n .* A matrix A is positive (semi)definite if and only if (i) the eigenvalues of A are positive (non-negative), (ii) the principal minors of A are positive (non-negative), or (iii) the principal invariants of A are positive (non-negative). The principal minors of A are determinants of the principal submatrices, which are obtained by deleting k rows and columns from A in a symmetric way, i.e. both the i_1, \dots, i_k rows and the i_1, \dots, i_k columns are deleted. The largest principal submatrix of A is A itself.

On the other hand, copositive matrices are demanded to be positive not for all vectors in the \mathbb{R}_n , but only on positive vectors in \mathbb{R}_+^n . *A symmetric matrix A is copositive if the quadratic form $x^T A x \geq 0$ for all vectors $x \geq 0$ in the non-negative orthant \mathbb{R}_+^n .* (The notation $x \geq 0$ means that $x_i \geq 0$ for each $i = 0, \dots, n$.) *A symmetric matrix A is strictly copositive if the quadratic form $x^T A x > 0$ for all vectors $x > 0$ in the non-negative orthant \mathbb{R}_+^n .*

For matrices of low order the copositivity conditions are relatively simple. A symmetric matrix A of order 2 is copositive if and only if [25]

$$a_{11} \geq 0, \quad a_{22} \geq 0, \quad a_{12} + \sqrt{a_{11}a_{22}} \geq 0. \quad (6)$$

A symmetric matrix A of order 3 is copositive if and only if [26, 27]

$$\begin{aligned} a_{11} &\geq 0, & a_{22} &\geq 0, & a_{33} &\geq 0, \\ \bar{a}_{12} = a_{12} + \sqrt{a_{11}a_{22}} &\geq 0, & \bar{a}_{13} = a_{13} + \sqrt{a_{11}a_{33}} &\geq 0, & \bar{a}_{23} = a_{23} + \sqrt{a_{22}a_{33}} &\geq 0, \\ \sqrt{a_{11}a_{22}a_{33}} + a_{12}\sqrt{a_{33}} + a_{13}\sqrt{a_{22}} + a_{23}\sqrt{a_{11}} + \sqrt{2\bar{a}_{12}\bar{a}_{13}\bar{a}_{23}} &\geq 0. \end{aligned} \quad (7)$$

The Cottle-Habetler-Lemke theorem [28] provides a practical way to find analytical copositivity conditions for matrices of low order. *Let the order $n - 1$ submatrices of a real symmetric matrix A of order n be copositive. Then A is not copositive if and only if*

$$\det A < 0 \quad \wedge \quad \text{adj } A \geq 0, \quad (8)$$

or A is copositive if and only if

$$\det A \geq 0 \quad \vee \quad (\text{adj } A)_{ij} < 0 \text{ for some } i, j. \quad (9)$$

The adjugate of A is the transpose of the cofactor matrix of A :

$$(\text{adj } A)_{ij} = (-1)^{i+j} M_{ji}, \quad (10)$$

where M_{ij} is the (i, j) minor of A , the determinant of the submatrix resulting from deleting the i th row and j th column of A .

Another general way to test copositivity is Kaplan's test [29]: *A symmetric matrix A is copositive if and only if every principal submatrix of A has no eigenvector $v \geq 0$ with associated eigenvalue $\lambda \leq 0$.*

We see that while the positivity of the matrix can be checked via its eigenvalues and the direction of its eigenvectors is irrelevant in this case, to check for copositivity at worst all the eigenvalues and eigenvectors of all principal submatrices of the matrix have to be calculated.

Copositivity with respect to a closed cone K or K -copositivity means

$$x^T A x \geq 0 \quad \forall x \in K. \quad (11)$$

If the cone K is polyhedral, it can be represented in the form $K = \{Gx \mid x \in \mathbb{R}_+^p\}$, where G is a real matrix whose columns $\{u_1, \dots, u_n\}$ are positively linearly independent vectors in \mathbb{R}^n that map the extremal rays of \mathbb{R}_+^p (basis vectors) into the extremal rays of the cone K . In such a case, the condition (11) takes the form [5]

$$x^T G^T A G x \geq 0 \quad \forall x \in \mathbb{R}_+^p. \quad (12)$$

Therefore to check whether A is copositive on K , we can check the usual copositivity of $G^T A G$ on \mathbb{R}_+^p .

Kaplan's test can also be generalised to copositivity on a closed cone K by requiring that every principal submatrix of A have no eigenvector $v \in K$ with associated eigenvalue $\lambda \leq 0$.

2.2 Orbit Spaces

Because the scalar potential is a homogenous polynomial of fields, the question whether the potential is bounded from below does not depend on the norm of the field for a single scalar field (e.g. the Higgs in the SM) and depends only on ratios of the norms for multiple fields. Orbit spaces for different gauge groups and potentials were actively studied in the 1980s, especially in the context of Grand Unified Theories. We give a short review with references to some works. The space of magnitudes of fields is the non-negative orthant \mathbb{R}_+^n and hence the problem of positivity of the potential can be put in terms of copositivity of the matrix (or tensor) of quartic couplings on the basis of norms of fields, with a few orbit space parameters to minimise over.

Orbit spaces in the context of spontaneous symmetry breaking are described in [30, 31]. The method of orbit spaces to minimise potentials is detailed in [11]. The case of the adjoint representation of $SU(N)$, in particular of the **24** of $SU(5)$, is detailed in [10]. An illustration for a potential of two scalars in different representations is given in [9] for the orbit space for the quartic potential of the **5** and the **24** representation of $SU(5)$. In [12] the case of $SO(N)$ with adjoint + vector representation is analysed. For several groups, the orbit spaces for simple potentials are described in [13].

We present a short outline of the method for a potential of one scalar multiplet. In our cursory review we will heed to the presentation of Jai Sam Kim [11]. For a theory with the non-Abelian gauge group G , the quartic potential for a scalar ϕ in an n -dimensional irreducible representation **R** of G can be written as

$$V(\phi) = \lambda_\phi (\phi_i^* \phi_i)^2 + \lambda'_\phi f_{ijkl} \phi_i^* \phi_j \phi_k^* \phi_l + \lambda''_\phi g_{ijkl} \phi_i^* \phi_j \phi_k^* \phi_l + \dots, \quad (13)$$

where f and g specify different gauge invariant contractions of indices. The potential $V(\phi)$ is invariant under a group transformation

$$\phi_j = T(\theta)_{ji} \phi_i, \quad (14)$$

where $T(\theta)$ is an n -dimensional matrix representing a group element. In general

$$T(\theta) = e^{-i\theta_L X_L}, \quad (15)$$

where X_L are the generators of the group G and θ_L are parameters that specify the group element.

Because in general the multiplet ϕ has many components, it is hard to solve the minimisation equations (4) for them. Moreover, there is a degeneracy of the components of ϕ that give the same minimum of V . A gauge transformation rotates the components of ϕ while leaving the value of the potential unchanged.

More formally, the orbit of a particular ϕ with constant components (such as a vacuum expectation value) is the set of states $\phi_\theta = T(\theta)\phi$ with $T(\theta)$ an element of G . It can be shown that all the states ϕ_θ respect the same group, the little group of the orbit, as ϕ does. If the group is unitary, then all the states ϕ_θ have the same norm $\phi^\dagger\phi$. The set of orbits that respect the same little group is called the stratum of the little group. Therefore we have to look for the orbit – and its little group – that minimises the potential.

Orbits of ϕ are specified by invariant polynomials $P(\phi)$ [32, 33, 34]. There is a basis set of invariant polynomials $I_a(\phi)$ such that every invariant polynomial $P(\phi)$ can be expressed as a polynomial in the polynomial basis: $P(\phi) = \bar{P}[I_a(\phi)]$. Each representation \mathbf{R} has a different number ℓ of basic invariants. An orbit can be pictured as a point in the ℓ -dimensional space of I_a .

The magnitude of ϕ is irrelevant to minimising the potential to find the vacuum stability conditions. The strata can be specified by dimensionless ratios of invariants, e.g.

$$\alpha_1 = \frac{f_{ijkl}\phi_i^*\phi_j\phi_k^*\phi_l}{(\phi_i^*\phi_i)^2}. \quad (16)$$

These dimensionless ratios are called orbit parameters and can be thought of as a set of angles. The potential (13) can then be written as

$$V(\phi) = \left[\lambda_\phi + \lambda'_\phi \alpha_1(\hat{\phi}) + \lambda''_\phi \alpha_2(\hat{\phi}) + \dots \right] |\phi|^4, \quad (17)$$

where

$$|\phi|^2 = \phi_i^*\phi_i, \quad \hat{\phi}_i = \frac{\phi_i}{|\phi|}. \quad (18)$$

The potential $V(\phi)$ is bounded from below if

$$\lambda_\phi + \lambda'_\phi \alpha_1(\hat{\phi}) + \lambda''_\phi \alpha_2(\hat{\phi}) + \dots > 0 \quad \text{for any } \alpha_i(\hat{\phi}). \quad (19)$$

Therefore, we have to minimise the potential with respect to the orbit space parameters. It is obvious that for any $\hat{\phi}$, the range of α_i is bounded from below and above: $\alpha_{i\min} \leq \alpha_i \leq \alpha_{i\max}$. We have to calculate the orbit space – the physical region in the orbit space parameters α_i . Because $\lambda_\phi + \lambda'_\phi \alpha_1 + \lambda''_\phi \alpha_2 = C$ describes a line in the orbit space, the minimum of the potential is on the boundary of the orbit space, in particular it can be on a cusp of the boundary curve.

Similar considerations apply for more than two orbit space parameters and for several scalars in different representations. For many more details, we refer the interested reader to the works cited.

3 Vacuum Stability of the Scalar Potential of Two Real Scalars

3.1 Vacuum Stability Conditions from Positivity of a Quartic Polynomial

The most general scalar potential of two real scalar fields ϕ_1 and ϕ_2 is⁵

$$V(\phi_1, \phi_2) = \lambda_{ij}\phi_1^i\phi_2^j = \lambda_{40}\phi_1^4 + \lambda_{31}\phi_1^3\phi_2 + \lambda_{22}\phi_1^2\phi_2^2 + \lambda_{13}\phi_1\phi_2^3 + \lambda_{04}\phi_2^4. \quad (20)$$

⁵Of course, the most general potential of the real and imaginary components of a complex singlet $S = \phi_1 + i\phi_2$ can be written in the same form. If $\lambda_{31} = \lambda_{13} = 0$, the potential has a CP symmetry. Therefore λ_{31} and λ_{13} could be naturally small.

Here we use the same notation λ_{ij} in a different way than in eq. (5) and copositivity, but the meaning should be clear from the context.

Set either ϕ_1 or ϕ_2 to zero, and it follows that the self-couplings λ_{40} and λ_{04} have to be positive in order for V to be positive.⁶ When both fields are non-zero, we can divide the potential V by ϕ_2^4 and choose the ratio $x = \phi_1/\phi_2$ as the new variable (equivalently, set $\phi_2 = 1$ to dehomogenise V), reducing the question of vacuum stability of the potential to that of the positivity of a general quartic polynomial of one variable,

$$P = a_4 x^4 + a_3 x^3 + a_2 x^2 + a_1 x + a_0, \quad (21)$$

which obviously is positive if it has no real roots and $a_4 > 0$ and $a_0 > 0$.

The nature of the roots of P can be determined by considering its discriminant

$$\begin{aligned} D = & 256a_0^3a_4^3 - 4a_1^3a_3^3 - 27a_0^2a_3^4 + 16a_0a_2^4a_4 - 6a_0a_1^2a_3^2a_4 - 27a_1^4a_4^2 \\ & - 192a_0^2a_1a_3a_4^2 - 4a_2^3(a_0a_3^2 + a_1^2a_4) + 18a_2(a_1a_3 + 8a_0a_4)(a_0a_3^2 + a_1^2a_4) \\ & + a_2^2(a_1^2a_3^2 - 80a_0a_1a_3a_4 - 128a_0^2a_4^2), \end{aligned} \quad (22)$$

and two additional polynomials of its coefficients,

$$Q = 8a_2a_4 - 3a_3^2, \quad R = 64a_0a_4^3 + 16a_2a_3^2a_4 - 16a_4^2(a_2^2 + a_1a_3) - 3a_3^4. \quad (23)$$

The condition for P to have only complex roots is [35, 36]

$$D > 0 \wedge (Q > 0 \vee R > 0). \quad (24)$$

In the marginal case $D = 0$ (which we can usually ignore), the conditions

$$Q > 0, \quad R = 0, \quad S = a_3^3 + 8a_1a_4^2 - 4a_4a_3a_2 = 0 \quad (25)$$

must hold.

For $V(\phi_1, \phi_2)/\phi_2^4$, the discriminant D and the polynomials Q and R are given by

$$\begin{aligned} D = & 256\lambda_{40}^3\lambda_{04}^3 - 4\lambda_{31}^3\lambda_{13}^3 - 27\lambda_{31}^4\lambda_{04}^2 + 16\lambda_{40}\lambda_{22}^4\lambda_{04} - 6\lambda_{40}\lambda_{31}^2\lambda_{04}\lambda_{13}^2 - 27\lambda_{40}^2\lambda_{13}^4 \\ & - 192\lambda_{40}^2\lambda_{31}\lambda_{04}^2\lambda_{13} - 4\lambda_{22}^3(\lambda_{31}^2\lambda_{04} + \lambda_{40}\lambda_{13}^2) + 18\lambda_{22}(8\lambda_{40}\lambda_{04} + \lambda_{31}\lambda_{13}) \\ & \times (\lambda_{31}^2\lambda_{04} + \lambda_{40}\lambda_{13}^2) + \lambda_{22}^2(\lambda_{31}^2\lambda_{13}^2 - 80\lambda_{40}\lambda_{31}\lambda_{04}\lambda_{13} - 128\lambda_{40}^2\lambda_{04}^2), \end{aligned} \quad (26)$$

$$Q = 8\lambda_{40}\lambda_{22} - 3\lambda_{31}^2, \quad (27)$$

$$R = 64\lambda_{40}^3\lambda_{04} + 16\lambda_{40}\lambda_{22}\lambda_{31}^2 - 16\lambda_{40}^2(\lambda_{22}^2 + \lambda_{31}\lambda_{13}) - 3\lambda_{31}^4, \quad (28)$$

and the vacuum stability conditions for $V(\phi_1, \phi_2)$ are given by

$$\lambda_{40} > 0, \quad \lambda_{04} > 0, \quad D > 0 \wedge (Q > 0 \vee R > 0). \quad (29)$$

For the record, the polynomial $S = \lambda_{13}^3 + 4\lambda_{40}(2\lambda_{40}\lambda_{31} - \lambda_{22}\lambda_{13})$.

Of course, we could as well divide by ϕ_1^4 and choose ϕ_2/ϕ_1 as the variable.⁷ The discriminant D is invariant under the exchange, but Q and R are not. The allowed parameter space, of course, stays the same. If either λ_{31} or λ_{13} is zero due to some symmetry, this freedom permits us to simplify the expressions for Q and R . For example, if $\lambda_{31} = 0$, the conditions are simpler if we choose ϕ_1/ϕ_2 as the

⁶Note that we can scale the fields by $\phi_1 \rightarrow \phi_1/\lambda_{40}^{1/4}$ and $\phi_2 \rightarrow \phi_2/\lambda_{04}^{1/4}$, so in effect there are three independent parameters to consider.

⁷In fact, it is possible to not only exchange the fields but to rotate them by an arbitrary angle α .




Figure 1: Left panel: An example of parameter space allowed by vacuum stability constraints (29) for the potential (20) two real scalars (dark green). If the scalars take only non-negative values, the light green area is allowed in addition (32). The values of the remaining parameters are $\lambda_{40} = 0.125$, $\lambda_{04} = 0.25$ and $\lambda_{22} = 0.25$. Right panel: The allowed parameter space (dark green) in the λ_{22} vs. λ_{40} plane with $\lambda_{04} = 0.25$, $\lambda_{13} = -0.75$ and $\lambda_{31} = 0$. The dashed line in the right panel shows the vacuum stability bound (31) from copositivity for $\lambda_{31} = \lambda_{13} = 0$.

variable, since the remaining λ_{13} term is only linear in ϕ_1 . In this case the condition $Q > 0 \vee R > 0$ takes the form $\lambda_{22} + 2\sqrt{\lambda_{40}\lambda_{04}} > 0$.

As a cross-check, we can set both λ_{31} and λ_{13} to zero. Then the vacuum stability conditions become

$$\begin{aligned} \lambda_{40} > 0, \quad \lambda_{04} > 0, \quad D = \lambda_{40}\lambda_{04}(\lambda_{22}^2 - 4\lambda_{40}\lambda_{04})^2 > 0, \\ Q = \lambda_{40}^2(4\lambda_{40}\lambda_{04} - \lambda_{22}^2) > 0 \vee R = \lambda_{40}\lambda_{22} > 0, \end{aligned} \quad (30)$$

which can be simplified to

$$\lambda_{40} > 0, \quad \lambda_{04} > 0, \quad \lambda_{22} + 2\sqrt{\lambda_{40}\lambda_{04}} > 0, \quad (31)$$

the usual copositivity conditions for V with ϕ_1^2 and ϕ_2^2 as the non-negative variables.

An illustration of the vacuum stability conditions for some values of parameters is given in Figure 1. The parts of the parameter space that are allowed are shown in dark green. In addition, the light green area is allowed if the scalars can have only non-negative values (see Section 3.2). If both $\lambda_{31} = \lambda_{13} = 0$, the vacuum stability conditions reduce to the usual copositivity conditions (31) (dashed line in the right panel).

3.2 Bounded From Below Conditions for a Quartic Polynomial on \mathbb{R}_+

The domain of the general quartic polynomial (21) can be restricted to *non-negative* real numbers \mathbb{R}_+ . In this case, the allowed range of parameters is somewhat larger. The positivity conditions for the polynomial

(21) for $x \geq 0$ are given by [37]⁸

$$\begin{aligned} & (D \leq 0 \wedge a_3 \sqrt{a_0} + a_1 \sqrt{a_4} > 0) \\ & \vee (-2\sqrt{a_0 a_4} < a_2 < 6\sqrt{a_0 a_4} \wedge D \geq 0 \wedge \Lambda_1 \leq 0) \\ & \vee (6\sqrt{a_0 a_4} < a_2 \wedge [(a_1 > 0 \wedge a_3 > 0) \vee (D \geq 0 \wedge \Lambda_2 \leq 0)]), \end{aligned} \quad (32)$$

where

$$\Lambda_1 = (\sqrt{a_0} a_3 - a_1 \sqrt{a_4})^2 - 32(a_0 a_4)^{\frac{3}{2}} - 16 \left(a_0 a_2 a_4 + a_0^{\frac{5}{4}} a_3 a_4^{\frac{3}{4}} + a_0^{\frac{3}{4}} a_1 a_4^{\frac{5}{4}} \right), \quad (33)$$

$$\Lambda_2 = (\sqrt{a_0} a_3 - a_1 \sqrt{a_4})^2 - \frac{4\sqrt{a_0 a_4}(a_2 + 2\sqrt{a_0 a_4})(\sqrt{a_0} a_3 + a_1 \sqrt{a_4} + 4\sqrt{a_0 a_4} \sqrt{a_2 - 2\sqrt{a_0 a_4}})}{\sqrt{a_2 - 2\sqrt{a_0 a_4}}}. \quad (34)$$

The main difference with the case of two real scalars is that a range of positive and opposite-sign λ_{31} and λ_{13} is allowed, as seen in the left hand panel of figure 1. Also note that it is trivial to restrict the domain to the negative numbers instead by taking $x \rightarrow -x$, equivalent to changing $a_1 \rightarrow -a_1$ and $a_3 \rightarrow -a_3$ in the above conditions.

If the non-negative variables are magnitudes of scalar fields, then the coefficients a_i may depend on additional orbit space parameters, notably phases. These may allow, in effect, to always choose the λ_{31} and/or λ_{13} terms to be negative. For that reason, as we will see below for the 2HDM, the conditions (32) for a positive variable can often be eschewed in favour of the simpler conditions (24) for a real variable.

3.3 Vacuum Stability Conditions from Positivity with an Affine Space

We will derive another, different form of vacuum stability conditions for the potential (20). The matrix of quartic couplings of the potential (20) in the monomial basis $(\phi_1^2, \phi_1 \phi_2, \phi_2^2)$ is

$$\Lambda = \begin{pmatrix} \lambda_{40} & \frac{1}{2}\lambda_{31} & \frac{1}{2}(1-c)\lambda_{22} \\ \frac{1}{2}\lambda_{31} & c\lambda_{22} & \frac{1}{2}\lambda_{13} \\ \frac{1}{2}(1-c)\lambda_{22} & \frac{1}{2}\lambda_{13} & \lambda_{04} \end{pmatrix}, \quad (35)$$

where c is an arbitrary constant due to the ambiguity $(\phi_1 \phi_2)^2 = \phi_1^2 \phi_2^2$. The matrices $\Lambda(c)$ form an affine space. If for some value of c the matrix Λ is positive-definite, then the potential (20) is bounded from below. Note that since $\phi_1 \phi_2 \in \mathbb{R}$, one has to demand the usual positivity, not copositivity.

The Sylvester criterion for the positivity of Λ is given by

$$\lambda_{40} > 0, \quad \lambda_{04} > 0, \quad c\lambda_{22} > 0, \quad (36)$$

$$4\lambda_{40}\lambda_{04} + (1-c)^2\lambda_{22}^2 > 0, \quad 4c\lambda_{22}\lambda_{40} - \lambda_{31}^2 > 0, \quad 4c\lambda_{22}\lambda_{04} - \lambda_{13}^2 > 0, \quad (37)$$

$$c\lambda_{22}(4\lambda_{40}\lambda_{04} - \lambda_{31}\lambda_{13}) - \lambda_{40}\lambda_{13}^2 + \lambda_{22}\lambda_{31}\lambda_{13} - \lambda_{04}\lambda_{31}^2 - c(1-c)^2\lambda_{22}^3 > 0. \quad (38)$$

First of all, the last inequality of (36) implies that c is real and has the same sign as λ_{22} . The first inequality of (37) is trivially satisfied. To satisfy the last two inequalities of (37) one must have $|c| \geq |c_0|$, where

$$c_0 \equiv \frac{1}{4\lambda_{22}} \max \left(\frac{\lambda_{13}^2}{\lambda_{40}}, \frac{\lambda_{31}^2}{\lambda_{04}} \right). \quad (39)$$

⁸We have restored $a \equiv a_0$ and $e \equiv a_4$ in the conditions given in [37] and slightly reorganised them. They were first used for finding vacuum stability conditions in [38].

The extrema of the left hand side (LHS) of the inequality (38) with respect to c are

$$c_{\pm} = \frac{1}{3} \left[2 \pm \frac{\sqrt{\lambda_{22}^2(\lambda_{22}^2 + 12\lambda_{40}\lambda_{04} - 3\lambda_{31}\lambda_{13})}}{\lambda_{22}^2} \right]. \quad (40)$$

Which of c_{\pm} gives the maximum of the LHS of (38)? The second derivative of the LHS of (38) with respect to c is $(1 - \frac{3}{2}c)\lambda_{22}^3$. Inserting here the solutions c_{\pm} , we see that for $\lambda_{22} > 0$, the maximum is given by c_+ and likewise for $\lambda_{22} < 0$ the maximum is c_- . Of course, if for $\lambda_{22} > 0$, for example, $c_+ < c_0$, one must take $c = c_0$.

Therefore, the optimal value for c is

$$c = \begin{cases} \max(c_0, c_+) & \text{if } \lambda_{22} > 0, \\ \min(c_0, c_-) & \text{if } \lambda_{22} < 0, \\ c = 0 & \text{if } \lambda_{22} = 0. \end{cases} \quad (41)$$

The conditions (36), (37) and (38) together with (41) define the same region of the parameter space as (29). We have traded the relative complexity of the latter for the apparent simplicity of the former at the cost of introducing the optimal coefficient c as a piece-wise function of the couplings that has a discontinuity at $\lambda_{22} = 0$.

4 Vacuum Stability for Two Real Scalars & the Higgs boson

The most general scalar potential of two real scalar fields ϕ_1 and ϕ_2 and the Higgs doublet H is

$$\begin{aligned} V(\phi_1, \phi_2, |H|^2) &= \lambda_H |H|^4 + \lambda_{H20} |H|^2 \phi_1^2 + \lambda_{H11} |H|^2 \phi_1 \phi_2 + \lambda_{H02} |H|^2 \phi_2^2 \\ &\quad + \lambda_{40} \phi_1^4 + \lambda_{31} \phi_1^3 \phi_2 + \lambda_{22} \phi_1^2 \phi_2^2 + \lambda_{13} \phi_1 \phi_2^3 + \lambda_{04} \phi_2^4 \\ &\equiv \lambda_H |H|^4 + M^2(\phi_1, \phi_2) |H|^2 + V(\phi_1, \phi_2), \end{aligned} \quad (42)$$

where $M^2(\phi_1, \phi_2) \equiv \lambda_{H20} \phi_1^2 + \lambda_{H11} \phi_1 \phi_2 + \lambda_{H02} \phi_2^2$ and $V(\phi_1, \phi_2) \equiv V(\phi_1, \phi_2, 0)$.

The potential (42) is a quadratic polynomial in $|H|^2$. Setting $\phi_1 = \phi_2 = 0$, we obtain $\lambda_H > 0$. Setting $|H|^2 = 0$ we recover the conditions (29) for $V(\phi_1, \phi_2) > 0$. If all three fields are non-zero, we can eliminate the Higgs: $V(\phi_1, \phi_2, |H|^2) > 0$ if either $M^2(\phi_1, \phi_2) > 0$, or else the discriminant of $V(\phi_1, \phi_2, |H|^2)$ with respect to $|H|^2$ is negative, that is $M^4(\phi_1, \phi_2) - 4\lambda_H V(\phi_1, \phi_2) < 0$.

An equivalent way to eliminate $|H|^2$ is to minimise the potential with respect to it:

$$0 = \frac{\partial V}{\partial |H|^2} = 2\lambda_H |H|^2 + M^2(\phi_1, \phi_2), \quad (43)$$

giving

$$|H|_{\min}^2 = -\frac{1}{2\lambda_H} M^2(\phi_1, \phi_2). \quad (44)$$

Again, either $M^2(\phi_1, \phi_2) > 0$ and the solution for $|H|_{\min}^2$ is unphysical, or else $V_{|H|^2=|H|_{\min}^2} = V(\phi_1, \phi_2) - \frac{1}{4\lambda_H} M^4(\phi_1, \phi_2)$ must be positive. Inserting the solution (44) into the potential in effect means substituting $\lambda_{40} \rightarrow \lambda_{40} - \lambda_{H20}^2/\lambda_H$ and so on in $V(\phi_1, \phi_2)$.

Therefore, for the potential (42) to be bounded from below, we altogether require

$$\lambda_H > 0, \quad V(\phi_1, \phi_2) > 0, \quad M^2(\phi_1, \phi_2) > 0 \vee V_{|H|^2=|H|_{\min}^2}(\phi_1, \phi_2) = V(\phi_1, \phi_2) - \frac{1}{4\lambda_H} M^4(\phi_1, \phi_2) > 0, \quad (45)$$

where the last OR condition is tantamount to the problem of positivity of the quartic polynomial $V_{|H|^2=|H|_{\min}^2}$ with the quadratic constraint $M^2 < 0$. In general $M^2(\phi_1, \phi_2)$ can be positive for some values of ϕ_1 and ϕ_2 and negative for others. In this case the region defined by $M^2(\phi_1, \phi_2) < 0$ is a pointed double cone in the $\phi_1\phi_2$ -plane.

In some regions of the parameter space it is easy to find the conditions. If the coefficient matrix of M^2 , given by

$$\mathbf{M}^2 = \begin{pmatrix} \lambda_{H20} & \frac{1}{2}\lambda_{H11} \\ \frac{1}{2}\lambda_{H11} & \lambda_{H02} \end{pmatrix}, \quad (46)$$

is positive-definite, that is

$$\lambda_{H20} > 0, \quad \lambda_{H02} > 0, \quad 4\lambda_{H20}\lambda_{H02} > \lambda_{H11}^2, \quad (47)$$

then $M^2 > 0$ for any values of the fields. If, on the other hand,

$$\lambda_{H20} \leq 0, \quad \lambda_{H02} \leq 0, \quad 4\lambda_{H20}\lambda_{H02} \leq \lambda_{H11}^2, \quad (48)$$

then $M^2 \leq 0$ for any values of the fields, and $V_{|H|^2=|H|_{\min}^2} > 0$ must hold for all values of the fields.

The intermediate situation, where for some values of the fields $M^2 < 0$ and for others $M^2 > 0$, occurs if the eigenvalues of \mathbf{M}^2 have opposite sign, that is, the determinant of \mathbf{M}^2 is negative:

$$4\lambda_{H20}\lambda_{H02} < \lambda_{H11}^2. \quad (49)$$

Then we can always make a transformation of the singlet fields to bring M^2 into the anti-diagonal form

$$M'^2 = \lambda'_{H11} \phi_1 \phi_2. \quad (50)$$

It is evident that such an M'^2 is negative in two opposite quadrants in the $\phi_1\phi_2$ -plane. If we dehomogenise $V_{|H|^2=|H|_{\min}^2}$ by taking e.g. $\phi_1 = 1$, we have to restrict the other field ϕ_2 to a half-axis to respect the constraint $M'^2 < 0$. But we already have conditions for a quartic polynomial to be positive (or negative) on a half-axis: the conditions (32) for a quartic in a positive variable.

To begin to bring M^2 into the form (50), first we diagonalise \mathbf{M}^2 by the orthogonal matrix

$$\mathbf{U}_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}, \quad (51)$$

where

$$\sin \theta = \frac{\lambda_{H11} \operatorname{sgn}(\lambda_{H02} - \lambda_{H20})}{\sqrt{\lambda_{H11}^2 + \left[\sqrt{(\lambda_{H20} - \lambda_{H02})^2} + \sqrt{(\lambda_{H20} - \lambda_{H02})^2 + \lambda_{H11}^2} \right]^2}}, \quad (52)$$

$$\cos \theta = \frac{\sqrt{(\lambda_{H20} - \lambda_{H02})^2} + \sqrt{(\lambda_{H20} - \lambda_{H02})^2 + \lambda_{H11}^2}}{\sqrt{\lambda_{H11}^2 + \left[\sqrt{(\lambda_{H20} - \lambda_{H02})^2} + \sqrt{(\lambda_{H20} - \lambda_{H02})^2 + \lambda_{H11}^2} \right]^2}}. \quad (53)$$

For our purposes, $\operatorname{sgn} 0 = 1$. The eigenvalues of \mathbf{M}^2 are given by

$$\lambda'_{H\mp} = \frac{1}{2} \left[\lambda_{H20} + \lambda_{H02} \mp \sqrt{\lambda_{H11}^2 + (\lambda_{H20} - \lambda_{H02})^2} \right]. \quad (54)$$

In the intermediate case, we have $\lambda'_{H-} < 0$ and $\lambda'_{H+} > 0$. After the diagonalisation

$$\mathbf{M}^2 = \begin{pmatrix} \lambda'_{H20} & 0 \\ 0 & \lambda'_{H02} \end{pmatrix} \quad (55)$$

with $|\lambda'_{H20}| = -\lambda'_{H-}$, $|\lambda'_{H02}| = \lambda'_{H+}$ if $\lambda_{H20} < \lambda_{H02}$, and $|\lambda'_{H20}| = \lambda'_{H+}$, $|\lambda'_{H02}| = -\lambda'_{H-}$ if $\lambda_{H20} \geq \lambda_{H02}$, so

$$|\lambda'_{H20,H02}| = \mp \frac{1}{2} \text{sgn}(\lambda_{H02} - \lambda_{H20}) \left[\lambda_{H20} + \lambda_{H02} \mp \text{sgn}(\lambda_{H02} - \lambda_{H20}) \sqrt{\lambda_{H11}^2 + (\lambda_{H20} - \lambda_{H02})^2} \right]. \quad (56)$$

Next we scale \mathbf{M}^2 by

$$\mathbf{S} = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{1}{\sqrt{|\lambda'_{H20}|}} & 0 \\ 0 & \frac{1}{\sqrt{|\lambda'_{H02}|}} \end{pmatrix} \quad (57)$$

to make it proportional to the unit matrix. Finally we rotate \mathbf{M}^2 by

$$\mathbf{U}_{\frac{\pi}{4}} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \quad (58)$$

into the anti-diagonal form. Altogether, we transform the fields in $V_{|H|^2=|H|_{\min}^2}$ by

$$\begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} \rightarrow \mathbf{U}_{\frac{\pi}{4}}^T \mathbf{S} \mathbf{U}_{\frac{\pi}{4}} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}, \quad (59)$$

yielding the transformed potential $V'_{|H|^2=|H|_{\min}^2}$.

After the transformation, the coefficient matrix has the form

$$\mathbf{M}'^2 = \frac{1}{2} \text{sgn}(\lambda_{H02} - \lambda_{H20}) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (60)$$

hence $\lambda'_{H11} = \text{sgn}(\lambda_{H02} - \lambda_{H20})$.

We can now dehomogenise $V'_{|H|^2=|H|_{\min}^2}$ by taking $\phi_1 = 1$, and use the conditions (32) for when M'^2 is negative. If $\lambda'_{H11} < 0$, then M'^2 is negative in the 1st and 3rd quadrants of the $\phi_1\phi_2$ -plane and the conditions (32) apply as is; if $\lambda'_{H11} > 0$, then M'^2 is negative in the 2nd and 4th quadrants, and we have to take $\phi_2 \rightarrow -\phi_2$ in $V'_{|H|^2=|H|_{\min}^2}$ before applying the conditions. In short, we have to take $\phi_2 \rightarrow -\text{sgn}(\lambda'_{H11}) \phi_2$, equivalent to taking $\lambda'_{31} \rightarrow -\text{sgn}(\lambda'_{H11}) \lambda'_{31}$ and $\lambda'_{13} \rightarrow -\text{sgn}(\lambda'_{H11}) \lambda'_{13}$.

Altogether, the vacuum stability conditions for the potential (42) are given by

$$\begin{aligned} & \lambda_{40} > 0, \quad \lambda_{04} > 0, \quad \lambda_H > 0, \quad D_{|H|^2=0} \wedge (Q_{|H|^2=0} > 0 \vee R_{|H|^2=0} > 0), \\ & \lambda_{H20} \leq 0 \wedge \lambda_{H02} \leq 0 \wedge \lambda_{H11}^2 \leq 4\lambda_{H20}\lambda_{H02} \implies 4\lambda_H\lambda_{40} - \lambda_{H20}^2 > 0 \wedge 4\lambda_H\lambda_{04} - \lambda_{H02}^2 > 0 \\ & \wedge D_{|H|^2=|H|_{\min}^2} \wedge (Q_{|H|^2=|H|_{\min}^2} > 0 \vee R_{|H|^2=|H|_{\min}^2} > 0), \\ & \lambda_{H11}^2 > 4\lambda_{H20}\lambda_{H02} \implies \lambda'_{04} > 0 \wedge \lambda'_{40} > 0 \\ & \wedge \left[(D_{|H|^2=|H|_{\min}^2} \leq 0 \wedge (\lambda'_{31}\sqrt{\lambda'_{04}} + \lambda'_{13}\sqrt{\lambda'_{40}}) > 0) \right. \\ & \vee \left(-2\sqrt{\lambda'_{04}\lambda'_{40}} < \lambda'_{22} < 6\sqrt{\lambda'_{04}\lambda'_{40}} \wedge D_{|H|^2=|H|_{\min}^2} \geq 0 \wedge \Lambda'_{1|H|^2=|H|_{\min}^2} \leq 0 \right) \\ & \left. \vee \left(6\sqrt{\lambda'_{04}\lambda'_{40}} < \lambda'_{22} \wedge [(\lambda'_{13} > 0 \wedge \lambda'_{31} > 0) \vee (D_{|H|^2=|H|_{\min}^2} \geq 0 \wedge \Lambda'_{2|H|^2=|H|_{\min}^2} \leq 0)] \right) \right], \end{aligned} \quad (61)$$




Figure 2: An example of parameter space allowed by the vacuum stability conditions (61) for the potential (42) of two real scalars and the Higgs. The values of parameters are $\lambda_H = 0.125$, $\lambda_{40} = 0.125$, $\lambda_{04} = 0.25$, $\lambda_{31} = \lambda_{13} = \lambda_{H20} = \lambda_{H02} = 0$, except for plots where these couplings vary. Both light and dark green regions are allowed for $\lambda_{H11} = 0$, while only the dark green region is allowed for $\lambda_{H11} = 0.5$.

where we take $\lambda'_{31} \rightarrow -\text{sgn}(\lambda_{H02} - \lambda_{H20})\lambda'_{31}$ and $\lambda'_{13} \rightarrow -\text{sgn}(\lambda_{H02} - \lambda_{H20})\lambda'_{13}$. The primes on Λ'_1 and Λ'_2 indicate that they are calculated for the transformed potential $V'_{|H|^2=|H|^2_{\min}}$. The discriminant of the transformed potential is, up to a positive constant coefficient which does not affect positivity, equal to the discriminant of $V_{|H|^2=|H|^2_{\min}}$ and does not depend on $\text{sgn}(\lambda_{H02} - \lambda_{H20})$. The conditions (61) are illustrated in Figure 2.

For comparison, if we set all the terms containing odd powers of ϕ_1 and ϕ_2 to zero, $\lambda_{H11} = \lambda_{31} = \lambda_{13} = 0$, then the positivity of the potential is given by the much simpler (strict) copositivity constraints on the matrix of couplings in the $(\phi_1, \phi_2, |H|^2)$ basis,

$$\begin{aligned} \lambda_{40} &> 0, & \lambda_{04} &> 0, & \lambda_H &> 0, \\ \bar{\lambda}_{22} \equiv \lambda_{22} + 2\sqrt{\lambda_{40}\lambda_{04}} &> 0, & \bar{\lambda}_{H20} \equiv \lambda_{H20} + 2\sqrt{\lambda_{40}\lambda_H} &> 0, & \bar{\lambda}_{H02} \equiv \lambda_{H02} + 2\sqrt{\lambda_{04}\lambda_H} &> 0, \\ \sqrt{\lambda_{40}\lambda_{H02}} + \sqrt{\lambda_{04}\lambda_{H20}} + \sqrt{\lambda_H\lambda_{22}} + \sqrt{\lambda_{40}\lambda_{04}\lambda_H} + \sqrt{\bar{\lambda}_{22}\bar{\lambda}_{H20}\bar{\lambda}_{H02}} &> 0, \end{aligned} \quad (62)$$

which are still necessary, but not sufficient conditions for positivity in the general case.

The transformed potential $V'_{|H|^2=|H|^2_{\min}}$ is rather complicated, so let us look at special cases. If $\lambda_{H20} = \lambda_{H02} = 0$, then \mathbf{M}^2 already has the anti-diagonal form, $V'_{|H|^2=|H|^2_{\min}} = V_{|H|^2=|H|^2_{\min}}$ and $\lambda'_{H11} = \lambda_{H11}$.

If, on the other hand, $\lambda_{H11} = 0$, then \mathbf{M}^2 is diagonal and U_θ is the unit matrix. Then λ_{H20} and λ_{H02} must have opposite signs for the determinant of \mathbf{M}^2 to be negative and $\text{sgn}(\lambda_{H02} - \lambda_{H20}) = \text{sgn} \lambda_{H02}$. The

transformed potential (multiplied by the irrelevant positive coefficient $4\lambda_{H20}^2\lambda_{H02}^2$ to make it simpler) is

$$4\lambda_{H20}^2\lambda_{H02}^2V'_{|H|^2=|H|_{\min}^2} = \lambda'_{40}\phi_1^4 + \lambda'_{31}\phi_1^3\phi_2 + \lambda'_{22}\phi_1^2\phi_2^2 + \lambda'_{13}\phi_1\phi_2^3 + \lambda'_{04}\phi_2^4, \quad (63)$$

where

$$\begin{aligned} \lambda'_{40} &= \lambda_H[\lambda_{H20}^2\lambda_{04} - \lambda_{22}\lambda_{H20}\lambda_{H02} + \lambda_{40}\lambda_{H02}^2 + \sqrt{-\lambda_{H20}\lambda_{H02}}(\lambda_{13}|\lambda_{H20}| + \lambda_{31}|\lambda_{H02}|)], \\ \lambda'_{31} &= 2\lambda_H[2\lambda_{H20}^2\lambda_{04} - 2\lambda_{40}\lambda_{H02}^2 + \sqrt{-\lambda_{H20}\lambda_{H02}}(\lambda_{13}|\lambda_{H20}| - \lambda_{31}|\lambda_{H02}|)], \\ \lambda'_{22} &= 2\lambda_H(3\lambda_{40}\lambda_{H02}^2 + \lambda_{22}\lambda_{H20}\lambda_{H02} + 3\lambda_{04}\lambda_{H20}^2) - 4\lambda_{H20}^2\lambda_{H02}^2, \\ \lambda'_{13} &= 2\lambda_H[2\lambda_{H20}^2\lambda_{04} - 2\lambda_{40}\lambda_{H02}^2 - \sqrt{-\lambda_{H20}\lambda_{H02}}(\lambda_{13}|\lambda_{H20}| - \lambda_{31}|\lambda_{H02}|)], \\ \lambda'_{04} &= \lambda_H[\lambda_{H20}^2\lambda_{04} - \lambda_{22}\lambda_{H20}\lambda_{H02} + \lambda_{40}\lambda_{H02}^2 - \sqrt{-\lambda_{H20}\lambda_{H02}}(\lambda_{13}|\lambda_{H20}| + \lambda_{31}|\lambda_{H02}|)]. \end{aligned} \quad (64)$$

In several particle physics models, the equivalent of M^2 already is diagonal. An example is given by the classically scale invariant \mathbb{Z}_3 symmetric dark matter model [38].

5 Vacuum Stability of the 2HDM with Real Couplings

The scalar potential of two Higgs doublets H_1 and H_2 in the 2HDM with no explicit CP-violation is

$$\begin{aligned} V &= \lambda_1|H_1|^4 + \lambda_2|H_2|^4 + \lambda_3|H_1|^2|H_2|^2 + \lambda_4(H_1^\dagger H_2)(H_2^\dagger H_1) + \frac{1}{2}\lambda_5[(H_1^\dagger H_2)^2 + (H_2^\dagger H_1)^2] \\ &\quad + \lambda_6|H_1|^2(H_1^\dagger H_2 + H_2^\dagger H_1) + \lambda_7|H_2|^2(H_1^\dagger H_2 + H_2^\dagger H_1) \\ &= \lambda_1 h_1^4 + \lambda_2 h_2^4 + \lambda_3 h_1^2 h_2^2 + \lambda_4 \rho^2 h_1^2 h_2^2 + \lambda_5 \rho^2 \cos 2\phi h_1^2 h_2^2 + 2\lambda_6 \rho \cos \phi h_1^3 h_2 + 2\lambda_7 \rho \cos \phi h_1 h_2^3, \end{aligned} \quad (65)$$

where we have taken the potentially complex couplings λ_5 , λ_6 and λ_7 real and parameterised the field bilinears as [39]

$$|H_1|^2 = h_1^2, \quad |H_2|^2 = h_2^2, \quad H_1^\dagger H_2 = h_1 h_2 \rho e^{i\phi}. \quad (66)$$

The orbit space parameter $\rho \in [0, 1]$ as implied by the Cauchy inequality $0 \leq |H_1^\dagger H_2| \leq |H_1||H_2|$. While the general form of the vacuum stability conditions for the most general 2HDM potential has been given [15, 16, 17] in the elegant ‘light cone’ formalism, the conditions can be given in a simple explicit and analytical form in terms of the potential couplings only in special cases.

If $\lambda_6 = \lambda_7 = 0$, we recover the vacuum stability conditions [40, 41, 42, 43, 39] for the inert doublet model

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_3 + 2\sqrt{\lambda_1\lambda_2} > 0, \quad (67)$$

$$\lambda_3 + \lambda_4 - |\lambda_5| + 2\sqrt{\lambda_1\lambda_2} > 0. \quad (68)$$

If $\lambda_6 \neq 0$ or $\lambda_7 \neq 0$, then the condition $\lambda_3 + \lambda_4 - \lambda_5 + 2\sqrt{\lambda_1\lambda_2} > 0$ is a necessary condition [24].

Finding the minimum of the general potential (65) is complicated because of its non-linear dependence on orbit parameters ρ and ϕ [12]. It is practically impossible to minimise the polynomials D , Λ_1 and Λ_2 in the conditions (32) with respect to these parameters. Instead, we minimise the potential with respect to ϕ , ρ , h_1 , and h_2 , with the fields lying on the circle $h_1^2 + h_2^2 = 1$, enforced by a Lagrange multiplier λ




Figure 3: The allowed region in the λ_7 vs. λ_6 plane for the 2HDM with no CP-violation is the intersection (dark green) of the regions with $\phi = 0$ and $\phi = \pi$ (light green). Other parameters have values $\lambda_1 = 0.125$, $\lambda_2 = 0.0625$ and $\lambda_3 = \lambda_4 = \lambda_5 = 0$.

as in eq. (3). The minimisation equations are

$$h_1 h_2 \rho (2\lambda_5 \rho h_1 h_2 \cos \phi + \lambda_6 h_1^2 + \lambda_7 h_2^2) \sin \phi = 0, \quad (69)$$

$$h_1 h_2 [(\lambda_4 + \lambda_5 \cos 2\phi) \rho h_1 h_2 + (\lambda_6 h_1^2 + \lambda_7 h_2^2) \cos \phi] = 0, \quad (70)$$

$$4\lambda_1 h_1^3 + 2[\lambda_3 + (\lambda_4 + \lambda_5 \cos 2\phi) \rho^2] h_1 h_2^2 + 6\lambda_6 \rho \cos \phi h_1^2 h_2 + 2\lambda_7 \rho \cos \phi h_2^3 = \lambda h_1, \quad (71)$$

$$4\lambda_2 h_2^3 + 2[\lambda_3 + (\lambda_4 + \lambda_5 \cos 2\phi) \rho^2] h_1^2 h_2 + 2\lambda_6 \rho \cos \phi h_1^3 + 6\lambda_7 \rho \cos \phi h_1 h_2^2 = \lambda h_2, \quad (72)$$

$$h_1^2 + h_2^2 = 1. \quad (73)$$

Eq. (69) reduces to $\sin \phi = 0$ which yields $\phi = 0$ and $\phi = \pi$ or $\cos \phi = \pm 1$. (There is another solution for ϕ that holds in the special case $\lambda_4 = \lambda_5$, but in this limit it gives the same solution for V_{\min} as we will obtain below.) The solutions for $h_1 = 0$, $h_2 = 0$ and $\rho = 0$ reproduce the conditions (67). While each of the two solutions gives a region of parameter space corresponding to the allowed region (32) for a positive variable, the allowed region is their *intersection* which is given by the allowed region (24) of parameter space for a *real* variable for $V_{\phi=0}$ or $V_{\phi=\pi}$, illustrated in figure 3.

The extremum solutions to the equations (70), (71), (72) and (73) are given by

$$\rho^2 = \frac{(\lambda_6 h_1^2 + \lambda_7 h_2^2)^2}{h_1^2 h_2^2 (\lambda_4 + \lambda_5)^2}, \quad (74)$$

$$h_1^2 = \frac{1}{2} \frac{(2\lambda_2 - \lambda_3)(\lambda_4 + \lambda_5) + 2\lambda_7(\lambda_6 - \lambda_7)}{(\lambda_1 + \lambda_2 - \lambda_3)(\lambda_4 + \lambda_5) - (\lambda_6 - \lambda_7)^2}, \quad (75)$$

$$h_2^2 = \frac{1}{2} \frac{(2\lambda_1 - \lambda_3)(\lambda_4 + \lambda_5) - 2\lambda_6(\lambda_6 - \lambda_7)}{(\lambda_1 + \lambda_2 - \lambda_3)(\lambda_4 + \lambda_5) - (\lambda_6 - \lambda_7)^2}, \quad (76)$$

$$V_{\min} = \frac{1}{4} \frac{(\lambda_4 + \lambda_5)(4\lambda_1 \lambda_2 - \lambda_3^2) - 4(\lambda_1 \lambda_7^2 + \lambda_2 \lambda_6^2 - \lambda_3 \lambda_6 \lambda_7)}{(\lambda_1 + \lambda_2 - \lambda_3)(\lambda_4 + \lambda_5) - (\lambda_6 - \lambda_7)^2}. \quad (77)$$

Note that the the solution for ρ^2 in (74) is non-negative if h_1^2 and h_2^2 are non-negative.




Figure 4: The allowed parameter space for some values of the couplings of the 2HDM. For all panels $\lambda_2 = 0.25$, $\lambda_4 = 0.25$, $\lambda_5 = 0$. The values of other parameters are $\lambda_1 = 0.25$, $\lambda_3 = 0$ for the first panel, $\lambda_1 = 0.25$, $\lambda_7 = 0.25$ for the second panel, and $\lambda_6 = 0.25$, $\lambda_7 = 0$ for the third panel (the dashed line shows the bound from copositivity for $\lambda_6 = \lambda_7 = 0$).

Because the extremum may not be a minimum or may lie outside the rectangular orbit space $\rho \in [0, 1]$, $\cos \phi \in [-1, 1]$, the potential must be separately minimised on the edges and vertices of that square. The potential is required to be positive in these parameter regions in any case as a necessary condition.

The conditions for $\rho = 0$ (and any ϕ) are given by (67). The extremum solutions (75), (76) and (74) are *already* calculated for $\cos \phi = \pm 1$. By continuity, the minimum value of $\cos \phi$ at $\rho = 1$ must be $\cos \phi = \pm 1$ as well. Thus the conditions for $\rho = 1$ can be found by applying the conditions (24) to $V_{\cos \phi = \pm 1, \rho = 1}/h_2^4$ with $x = h_1/h_2$. The discriminant D and the polynomials Q and R are the same for $\phi = 0$ and for $\phi = \pi$ and given by

$$D_{\cos \phi = \pm 1, \rho = 1} = 16[16\lambda_1^3\lambda_2^3 + \lambda_1\lambda_2\lambda_{345}^4 - 27\lambda_2^2\lambda_6^4 - 48\lambda_1^2\lambda_2^2\lambda_6\lambda_7 - 6\lambda_1\lambda_2\lambda_6^2\lambda_7^2 - 16\lambda_6^3\lambda_7^3 - 27\lambda_1^2\lambda_7^4 - \lambda_{345}^3(\lambda_2\lambda_6^2 + \lambda_1\lambda_7^2) + 18\lambda_{345}(2\lambda_1\lambda_2 + \lambda_6\lambda_7)(\lambda_2\lambda_6^2 + \lambda_1\lambda_7^2) + \lambda_{345}^2(-8\lambda_1^2\lambda_2^2 - 20\lambda_1\lambda_2\lambda_6\lambda_7 + \lambda_6^2\lambda_7^2)], \quad (78)$$

$$Q_{\cos \phi = \pm 1, \rho = 1} = 8\lambda_1\lambda_{345} - 12\lambda_6^2, \quad (79)$$

$$R_{\cos \phi = \pm 1, \rho = 1} = 16[4\lambda_1^3\lambda_2 + 4\lambda_1\lambda_{345}\lambda_6^2 - 3\lambda_6^4 - \lambda_1^2(\lambda_{345}^2 + 4\lambda_6\lambda_7)], \quad (80)$$

where $\lambda_{345} \equiv \lambda_3 + \lambda_4 + \lambda_5$.

Altogether, the conditions for the 2HDM potential with real couplings to be bounded from below are

$$V_{\rho=0} > 0 \wedge D_{\cos \phi = \pm 1, \rho = 1} \wedge (Q_{\cos \phi = \pm 1, \rho = 1} > 0 \vee R_{\cos \phi = \pm 1, \rho = 1} > 0) \wedge (0 < h_1^2 < 1 \wedge 0 < h_2^2 < 1 \wedge 0 < \rho^2 < 1 \implies V_{\min} > 0), \quad (81)$$

where the conditions for $V_{\rho=0} > 0$ are given by (67) and $p \implies q$ is equivalent to $\neg p \vee q$. In fact, it is enough to check that either h_1^2 or h_2^2 is within bounds, since they are related by $h_1^2 + h_2^2 = 1$. The conditions (81) are illustrated in Figure 4.

The approach we use gives a simpler result than directly minimising $V_{\cos \phi = \pm 1, \rho = 1}$ with respect to h_1 and h_2 as we did for V_{\min} , especially as the conditions (24) automatically take into account the two different values $\cos \phi = \pm 1$. For a potential that depends on three or more moduli of fields, e.g. the 3HDM, where we would have a h_3 besides h_1 and h_2 , this is not possible and one has to minimise the potential on a hypersphere or use tensor eigenvalues (see Section 7).

6 Vacuum Stability for \mathbb{Z}_3 Scalar Dark Matter

Another physical example is given by scalar dark matter stable under a \mathbb{Z}_3 discrete group. The most general scalar quartic potential of the SM Higgs H_1 , an inert doublet H_2 and a complex singlet S which is symmetric under a \mathbb{Z}_3 group is [22, 23]

$$\begin{aligned}
V(H_1, H_2, S) &= \lambda_1 |H_1|^4 + \lambda_2 |H_2|^4 + \lambda_3 |H_1|^2 |H_2|^2 + \lambda_4 (H_1^\dagger H_2)(H_2^\dagger H_1) + \lambda_S |S|^4 \\
&\quad + \lambda_{S1} |S|^2 |H_1|^2 + \lambda_{S2} |S|^2 |H_2|^2 + \frac{1}{2} (\lambda_{S12} S^2 H_1^\dagger H_2 + \lambda_{S12}^* S^{\dagger 2} H_2^\dagger H_1) \\
&= \lambda_1 h_1^4 + \lambda_2 h_2^4 + \lambda_3 h_1^2 h_2^2 + \lambda_4 \rho^2 h_1^2 h_2^2 + \lambda_S s^4 + \lambda_{S1} s^2 h_1^2 + \lambda_{S2} s^2 h_2^2 - |\lambda_{S12}| \rho s^2 h_1 h_2 \\
&\equiv \lambda_S s^4 + M^2(h_1, h_2) s^2 + V(h_1, h_2),
\end{aligned} \tag{82}$$

where we have used the parametrisation (66) for the doublet bilinears and $S = se^{i\phi_S}$, and we have minimised $\cos(\phi + 2\phi_S + \phi_{\lambda_{S12}}) = -1$ so $\lambda_{S12} = -|\lambda_{S12}|$ without loss of generality. We define $M^2(h_1, h_2) \equiv \lambda_{S1} h_1^2 - |\lambda_{S12}| \rho h_1 h_2 + \lambda_{S2} h_2^2$ and $V(h_1, h_2) \equiv V(h_1, h_2, 0)$.

The situation is similar to the case of two real scalars and the Higgs boson in Section 4. First of all, $\lambda_S > 0$ and $V(h_1, h_2) > 0$. The conditions for $V(h_1, h_2) > 0$ are the same as in the inert doublet model with $\lambda_5 = 0$:

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_3 + 2\sqrt{\lambda_1 \lambda_2} > 0, \tag{83}$$

$$\lambda_3 + \lambda_4 + 2\sqrt{\lambda_1 \lambda_2} > 0. \tag{84}$$

We minimise the potential with respect to h_1, h_2, s and ρ with the fields lying on a sphere, enforced by a Lagrange multiplier λ . The minimisation equations are

$$h_1 h_2 (2\rho \lambda_4 h_1 h_2 - |\lambda_{S12}| s^2) = 0, \tag{85}$$

$$4\lambda_1 h_1^3 + 2(\lambda_3 + \lambda_4 \rho^2) h_1 h_2^2 + 2\lambda_{S1} h_1 s^2 - |\lambda_{S12}| \rho h_2 s^2 = \lambda h_1, \tag{86}$$

$$4\lambda_2 h_2^3 + 2(\lambda_3 + \lambda_4 \rho^2) h_1^2 h_2 + 2\lambda_{S2} h_2 s^2 - |\lambda_{S12}| \rho h_1 s^2 = \lambda h_2, \tag{87}$$

$$s (4\lambda_S s^2 + 2\lambda_{S1} h_1^2 + 2\lambda_{S2} h_2^2 - 2|\lambda_{S12}| \rho h_1 h_2) = \lambda s, \tag{88}$$

$$h_1^2 + h_2^2 + s^2 = 1. \tag{89}$$

The solution with all fields non-zero is

$$\rho = \left(|\lambda_{S12}| s^2 \right) / \left(2\lambda_4 h_1 h_2 \right), \tag{90}$$

$$h_1^2 = \frac{1}{2} \left((2\lambda_2 - \lambda_3)(4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + 2\lambda_4 [(\lambda_3 + \lambda_{S1})\lambda_{S2} - 2\lambda_2 \lambda_{S1} - \lambda_{S2}^2] \right) / \left((\lambda_1 + \lambda_2 - \lambda_3) \right) \tag{91}$$

$$\times (4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + \lambda_4 [4\lambda_1 \lambda_2 - \lambda_3^2 - 4\lambda_1 \lambda_{S2} - 4\lambda_2 \lambda_{S1} + 2\lambda_3 (\lambda_{S1} + \lambda_{S2}) - (\lambda_{S1} - \lambda_{S2})^2],$$

$$h_2^2 = \frac{1}{2} \left((2\lambda_1 - \lambda_3)(4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + 2\lambda_4 [(\lambda_3 + \lambda_{S2})\lambda_{S1} - 2\lambda_1 \lambda_{S2} - \lambda_{S1}^2] \right) / \left((\lambda_1 + \lambda_2 - \lambda_3) \right) \tag{92}$$

$$\times (4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + \lambda_4 [4\lambda_1 \lambda_2 - \lambda_3^2 - 4\lambda_1 \lambda_{S2} - 4\lambda_2 \lambda_{S1} + 2\lambda_3 (\lambda_{S1} + \lambda_{S2}) - (\lambda_{S1} - \lambda_{S2})^2],$$

$$s^2 = \lambda_4 \left(4\lambda_1 \lambda_2 - \lambda_3^2 - 2\lambda_1 \lambda_{S2} - 2\lambda_2 \lambda_{S1} + \lambda_3 (\lambda_{S1} + \lambda_{S2}) \right) / \left((\lambda_1 + \lambda_2 - \lambda_3) \right) \tag{93}$$

$$\times (4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + \lambda_4 [4\lambda_1 \lambda_2 - \lambda_3^2 - 4\lambda_1 \lambda_{S2} - 4\lambda_2 \lambda_{S1} + 2\lambda_3 (\lambda_{S1} + \lambda_{S2}) - (\lambda_{S1} - \lambda_{S2})^2],$$

$$V_{\min} = \frac{1}{4} \left((4\lambda_1 \lambda_2 - \lambda_3^2)(4\lambda_S \lambda_4 - |\lambda_{S12}|^2) - 4\lambda_4 (\lambda_1 \lambda_{S2}^2 + \lambda_2 \lambda_{S1}^2 - \lambda_3 \lambda_{S1} \lambda_{S2}) \right) / \left((\lambda_1 + \lambda_2 - \lambda_3) \right) \tag{94}$$

$$\times (4\lambda_S \lambda_4 - |\lambda_{S12}|^2) + \lambda_4 [4\lambda_1 \lambda_2 - \lambda_3^2 - 4\lambda_1 \lambda_{S2} - 4\lambda_2 \lambda_{S1} + 2\lambda_3 (\lambda_{S1} + \lambda_{S2}) - (\lambda_{S1} - \lambda_{S2})^2].$$

Note that h_1^2 , h_2^2 , s^2 and V_{\min} share the same denominator. The solution $s = 0$ will repeat (83) and (84) and the solutions $h_1 = 0$ or $h_2 = 0$ will be made redundant by eq. (95).

Since $\rho = 0$ sets the λ_{S12} term to zero, $V_{\rho=0} > 0$ is biquadratic in the fields and we can calculate the positivity conditions for $V_{\rho=0} > 0$ via (strict) copositivity of the matrix of couplings in the (h_1^2, h_2^2, s^2) basis,

$$\begin{aligned} \lambda_S &> 0, & \lambda_1 &> 0, & \lambda_2 &> 0, \\ \bar{\lambda}_3 \equiv \lambda_3 + 2\sqrt{\lambda_1\lambda_2} &> 0, & \bar{\lambda}_{S1} \equiv \lambda_{S1} + 2\sqrt{\lambda_S\lambda_1} &> 0, & \bar{\lambda}_{S2} \equiv \lambda_{S2} + 2\sqrt{\lambda_S\lambda_2} &> 0, \\ \sqrt{\lambda_S\lambda_3} + \sqrt{\lambda_1\lambda_{S2}} + \sqrt{\lambda_2\lambda_{S1}} + \sqrt{\lambda_S\lambda_1\lambda_2} + \sqrt{\bar{\lambda}_{S1}\bar{\lambda}_{S2}\bar{\lambda}_3} &> 0, \end{aligned} \quad (95)$$

part of which repeats conditions (83).

For $\rho = 1$, instead of direct minimisation of the potential in all variables, we find it easier to calculate conditions very similar to the case (61) of two real scalars and the Higgs boson. To reduce $V_{\rho=1}$ to a polynomial of two variables, we minimise it with respect to s^2 :

$$s_{\min}^2 = -\frac{1}{2\lambda_S}M^2(h_1, h_2). \quad (96)$$

Again, either $M^2(h_1, h_2) > 0$ and the solution for s_{\min}^2 is unphysical, or else $V_{s^2=s_{\min}^2} = V(h_1, h_2) - \frac{1}{4\lambda_S}M^4(h_1, h_2) > 0$. We have

$$\begin{aligned} 4\lambda_S V_{s^2=s_{\min}^2} &= (4\lambda_1\lambda_S - \lambda_{S1}^2)h_1^4 + (4\lambda_2\lambda_S - \lambda_{S2}^2)h_2^4 + [4(\lambda_3 + \lambda_4)\lambda_S - 2\lambda_{S1}\lambda_{S2} - |\lambda_{S12}|^2]h_1^2h_2^2 \\ &\quad + \lambda_{S1}|\lambda_{S12}|h_1^3h_2 + \lambda_{S2}|\lambda_{S12}|h_1h_2^3. \end{aligned} \quad (97)$$

The coefficient matrix of M^2 is given by

$$\mathbf{M}^2 = \begin{pmatrix} \lambda_{S1} & -\frac{1}{2}|\lambda_{S12}| \\ -\frac{1}{2}|\lambda_{S12}| & \lambda_{S2} \end{pmatrix}. \quad (98)$$

Repeating the procedure of Section 4 step by step, the conditions for $V_{\rho=1} > 0$, in addition to $\lambda_S > 0$ and (83) and (84), are given by

$$\begin{aligned} \lambda_{S1} \leq 0 \wedge \lambda_{S2} \leq 0 \wedge |\lambda_{S12}|^2 \leq 4\lambda_{S1}\lambda_{S2} &\implies 4\lambda_S\lambda_1 - \lambda_{S1}^2 > 0 \wedge 4\lambda_S\lambda_2 - \lambda_{S2}^2 > 0 \\ \wedge D_{s^2=s_{\min}^2} \wedge (Q_{s^2=s_{\min}^2} > 0 \vee R_{s^2=s_{\min}^2} > 0), \\ |\lambda_{S12}|^2 > 4\lambda_{S1}\lambda_{S2} &\implies \lambda'_{40} > 0 \wedge \lambda'_{04} > 0 \\ \wedge \left[\left(D_{s^2=s_{\min}^2} \leq 0 \wedge \left(\lambda'_{31}\sqrt{\lambda'_{04}} + \lambda'_{13}\sqrt{\lambda'_{40}} \right) > 0 \right) \right. \\ \vee \left(-2\sqrt{\lambda'_{40}\lambda'_{04}} < \lambda'_{22} < 6\sqrt{\lambda'_{40}\lambda'_{04}} \wedge D_{s^2=s_{\min}^2} \geq 0 \wedge \Lambda'_{1\ s^2=s_{\min}^2} \leq 0 \right) \\ \left. \vee \left(6\sqrt{\lambda'_{04}\lambda'_{40}} < \lambda'_{22} \wedge [(\lambda'_{13} > 0 \wedge \lambda'_{31} > 0) \vee (D_{s^2=s_{\min}^2} \geq 0 \wedge \Lambda'_{2\ s^2=s_{\min}^2} \leq 0)] \right) \right], \end{aligned} \quad (99)$$

where we take $\lambda'_{31} \rightarrow -\text{sgn}(\lambda_{S2} - \lambda_{S1})\lambda'_{31}$ and $\lambda'_{13} \rightarrow -\text{sgn}(\lambda_{S2} - \lambda_{S1})\lambda'_{13}$. As in Section 4, λ'_{ij} denote the coefficients of the transformed potential $V'_{s^2=s_{\min}^2}$, where M'^2 is anti-diagonal.⁹ Note that we are justified to use the conditions (24) for the positivity of the general quartic on reals in the second line of (99), because in (97) the coefficients of the $h_1^3h_2$ and $h_1h_2^3$ terms are non-positive in that case.

⁹In eq. (99), we have retained the λ'_{ij} notation of Section 4 for ease of comparison. In the 2HDM notation, the coefficients would be $\lambda'_{40} \equiv \lambda'_1$, $\lambda'_{04} \equiv \lambda'_2$, $\lambda'_{22} \equiv \lambda'_{345}$, $\lambda'_{31} \equiv \lambda'_6$ and $\lambda'_{13} \equiv \lambda'_7$.




Figure 5: An example of parameter space allowed by the vacuum stability conditions (100) for the potential (82) of the SM Higgs, an inert doublet and a complex singlet. The values of parameters are $\lambda_H = 0.125$, $\lambda_S = 0.125$, $\lambda_2 = 0.25$, $\lambda_3 = \lambda_4 = \lambda_{S1} = \lambda_{S2} = 0$, except for plots where these couplings vary. Both light and dark green regions are allowed for for $|\lambda_{S12}| = 0$, while $|\lambda_{S12}| = 0.5$ only allows the dark green region. In the last two panels, where $|\lambda_{S12}|$ varies, only the dark blue region is allowed for $\lambda_{S2} = 0$, while $\lambda_{S2} = 0.5$ allows both the dark and light blue regions.

Altogether, the conditions for the potential (82) symmetric under a \mathbb{Z}_3 to be bounded from below are

$$\begin{aligned} &\lambda_S > 0 \wedge V(h_1, h_2) > 0 \wedge V_{\rho=0} > 0 \wedge V_{\rho=1} > 0 \\ &\wedge (0 < h_1^2 < 1 \wedge 0 < h_2^2 < 1 \wedge 0 < s^2 < 1 \wedge 0 < \rho^2 < 1 \implies V_{\min} > 0), \end{aligned} \quad (100)$$

where the conditions for $V(h_1, h_2) > 0$ are given by (83) and (84), the conditions for $V_{\rho=0} > 0$ are given by (95), the conditions for $V_{\rho=1} > 0$ are given by (99), and the extremum solutions h_1^2 , h_2^2 , s^2 , ρ^2 and V_{\min} are given by (90), (91), (92), (93) and (94). The conditions (100) are illustrated in Figure 5.

7 Tensor Eigenvalues

7.1 Positive Tensors

The most general scalar potential of n real singlet scalar fields ϕ_i can be written as

$$V = \lambda_{ijkl} \phi_i \phi_j \phi_k \phi_l \equiv \Lambda \phi^4, \quad (101)$$

where Λ is the tensor of scalar couplings and $\phi = (\phi_1, \dots, \phi_n)$. Clearly, the tensor Λ can always be made fully symmetric under permutations of the indices of its elements.

A symmetric matrix is positive if its eigenvalues are greater than zero. Can one generalise matrix eigenvalues and eigenvectors to tensors in such a way that they have similar properties? Indeed, tensor eigenvalues and eigenvectors have been defined independently by Qi in [44] (which we follow in our exposition) and Lim [45].

A real m th-order n -dimensional tensor A has n^m elements $A_{i_1 \dots i_m}$, where $i_j = 1, \dots, n$ for $j = 1, \dots, m$. A homogenous polynomial $f(x)$ of n variables and degree m can be written as the tensor product

$$f(x) = Ax^m \equiv A_{i_1 \dots i_m} x_{i_1} \cdots x_{i_m}. \quad (102)$$

In analogy with non-negative matrices, a n -dimensional tensor A is called non-negative if $Ax^m \geq 0$ for all $x \in \mathbb{R}^n$.

For a vector $x \in \mathbb{R}^n$ we define $(x^{[m]})_i = x_i^m$. The number λ is an eigenvalue of A if it is a solution to the equation

$$Ax^{m-1} = \lambda x^{[m-1]}, \quad (103)$$

where x is an eigenvector of A . For $m = 2$, the tensor A is a matrix and the equation (103) coincides with the usual matrix eigenvalue equation. When $m > 2$, the eigenvalues and eigenvectors of a tensor can be complex, but for even m , there always exist real eigenvalues and eigenvectors, so-called H -eigenvalues and H -eigenvectors.

An n -dimensional symmetric tensor of order m has $n(m-1)^{n-1}$ eigenvalues, so a coupling tensor of a renormalisable potential of n real scalar fields in 4-dimensional spacetime has $n3^{n-1}$ eigenvalues: for example, with two real fields, there are six eigenvalues. The product of all the eigenvalues of A is the hyperdeterminant $\det A$, that is the resultant of $Ax^{m-1} = 0$. When $m = 2$, the hyperdeterminant reduces to the usual matrix determinant. The sum of all eigenvalues is $(m-1)^{n-1}$ times the sum of diagonal elements or trace $\text{tr } A = \sum_i A_{iiii}$.

For even m , an eigenvalue equation similar to the matrix eigenvalue equation can be given,

$$\det(A - \lambda I) = 0, \quad (104)$$

where I is the unit tensor with $I_{i_1 \dots i_m} = \delta_{i_1 \dots i_m}$. The degree of the eigenvalue equation is $d = n(m-1)^{n-1}$.

Obviously only real eigenvectors have relevance to whether the tensor A is non-negative. *The tensor A is non-negative if all its eigenvalues with real eigenvectors are non-negative.* All the principal subtensors of A , that is the tensors obtained by setting one or more of the variables x_i in $f(x)$ to zero, must be positive as well.

Qi also defines tensor E -eigenvalues and E -eigenvectors, which are exactly the same as solutions to the equations (4) with fields constrained to a hypersphere. For even m , there always exist real E -eigenvalues and E -eigenvectors.

Copositive tensors are defined in obvious analogy to copositive matrices [46]: *A real symmetric tensor A of order m and dimension n is copositive if $Ax^m \geq 0$ for all $x \in \mathbb{R}_+^n$.* It has been shown that Kaplan's test of matrix co-positivity [29] directly generalises to copositive tensors [47]: *A symmetric tensor A is copositive if and only if every principal subtensor of A has no eigenvector $v > 0$ with associated H -eigenvalue $\lambda < 0$.*

7.2 Vacuum Stability of the Potential of Two Real Scalars

As an example, we consider again the general potential of two real scalars given by (20):

$$V(\phi_1, \phi_2) = \lambda_{ij} \phi_1^i \phi_2^j = \lambda_{40} \phi_1^4 + \lambda_{31} \phi_1^3 \phi_2 + \lambda_{22} \phi_1^2 \phi_2^2 + \lambda_{13} \phi_1 \phi_2^3 + \lambda_{04} \phi_2^4. \quad (105)$$

The tensor of the scalar couplings of the potential is given by

$$\Lambda = \left(\begin{pmatrix} \lambda_{40} & \frac{1}{4}\lambda_{31} \\ \frac{1}{4}\lambda_{31} & \frac{1}{6}\lambda_{22} \\ \frac{1}{4}\lambda_{31} & \frac{1}{6}\lambda_{22} \\ \frac{1}{6}\lambda_{22} & \frac{1}{4}\lambda_{13} \end{pmatrix} \begin{pmatrix} \frac{1}{4}\lambda_{31} & \frac{1}{6}\lambda_{22} \\ \frac{1}{6}\lambda_{22} & \frac{1}{4}\lambda_{13} \\ \frac{1}{6}\lambda_{22} & \frac{1}{4}\lambda_{13} \\ \frac{1}{4}\lambda_{13} & \lambda_{04} \end{pmatrix} \right), \quad (106)$$

that is, $\lambda_{1111} = \lambda_{40}$, $\lambda_{2222} = \lambda_{04}$, $\lambda_{1112} = \lambda_{1121} = \lambda_{1211} = \lambda_{2111} = \frac{1}{4}\lambda_{31}$ and so on.

The H -eigenvalue equations are

$$\begin{aligned} 4\lambda_{40}\phi_1^3 + 3\lambda_{31}\phi_1^2\phi_2 + 2\lambda_{22}\phi_1\phi_2^2 + \lambda_{13}\phi_2^3 &= 4\lambda\phi_1^3, \\ \lambda_{31}\phi_1^3 + 2\lambda_{22}\phi_1^2\phi_2 + 3\lambda_{13}\phi_1\phi_2^2 + 4\lambda_{04}\phi_2^3 &= 4\lambda\phi_2^3, \end{aligned} \quad (107)$$

The product of all solutions to the H -eigenvalue equations (107) is exactly the discriminant D given by (26) and their sum is $3(\lambda_{40} + \lambda_{04})$. The tensor eigenvalue equation (104) is a 6th degree equation, and the eigenvalues cannot in general be solved in radicals. Therefore, for two fields, we are better off using the conditions (29).

But for the general potential of three fields we have no such (relatively) simple conditions. For $m = 4$, $n = 3$, the recipe for the hyperdeterminant, given in [48], is very complicated in practice. Analytical expressions can be found only for coupling tensors of potentials that have few fields and are rather symmetric.

8 Conclusions

In particle physics, scalar potentials have to be bounded from below in order for the physics to make sense. Finding such conditions is a hard problem of algebraic geometry. We present analytical necessary and sufficient vacuum stability conditions for potentials of a few fields, where ‘few’ means two or more, depending on field content and symmetry. The vacuum stability conditions (29) for a general potential of two real fields fit on a few lines. Already for three fields, practical analytical conditions (61) can only be found for the potential (42), where at least one of the fields, such as the Higgs boson, is present only in biquadratic form. In this case the problem reduces to the positivity of a general quartic polynomial with a quadratic constraint.

As further examples that put several of the discussed techniques to use, we present simple vacuum stability conditions (81) for the 2HDM potential without explicit CP-violation, and vacuum stability conditions (100) for \mathbb{Z}_3 scalar dark matter with an inert doublet and a complex singlet. All analytical calculations have been checked numerically. We include a Mathematica notebook with the conditions with the \LaTeX source of the paper.

The vacuum stability conditions for the general potential of two real singlets (without or with the Higgs boson), and for the \mathbb{Z}_3 scalar dark matter are novel results. The vacuum stability conditions for the 2HDM potential with real couplings are in a shorter form than previous similar results [24].

Of course, our endeavour can be made much easier if a restrictive symmetry is imposed on the potential. If all the fields appear solely quadratically, for example, the problem becomes much simpler. Then the bounded from below conditions are given by copositivity constraints of the matrix of couplings. Many potentials can be written in terms of magnitudes of squares of fields and a few orbit space parameters.

The parameter space for more complicated potentials must be found numerically by minimising the potential on a hypersphere of field values or solving tensor eigenvalue equations. Still, some insight can be gained from (necessary) analytical conditions for a subspace where a field or more is set to zero, and the introduced methods can be used to reduce the parameter space for a numerical scan.

Acknowledgements

We would like to thank Hardi Veermäe for useful discussions. This work was supported by grants the Estonian Research Council grant PUT799, the grant IUT23-6 of the Estonian Ministry of Education and Research, and by the EU through the ERDF CoE program.

References

- [1] W. A. Bardeen, “On naturalness in the standard model.” FERMILAB-CONF-95-391-T, C95-08-27.3, 1995.
- [2] S. R. Coleman and E. J. Weinberg, *Radiative Corrections as the Origin of Spontaneous Symmetry Breaking*, *Phys.Rev.* **D7** (1973) 1888–1910.
- [3] K. Murty and S. Kabadi, *Some np-complete problems in quadratic and nonlinear programming*, *Mathematical Programming* **39** (1987) 117–129. 10.1007/BF02592948.
- [4] T. S. Motzkin, *Copositive quadratic forms*, in *National Bureau of Standards Report 1818*, pp. 11–22. 1952.
- [5] J.-B. Hiriart-Urruty and A. Seeger, *A variational approach to copositive matrices*, *SIAM Review* **52** (2010), no. 4 593–629.
- [6] K. Kannike, *Vacuum Stability Conditions From Copositivity Criteria*, *Eur.Phys.J.* **C72** (2012) 2093, [[arXiv:1205.3781](#)].
- [7] J.-B. Hiriart-Urruty and J. Malick, *A fresh variational-analysis look at the positive semidefinite matrices world*, *Journal of Optimization Theory and Applications* **153** (2012), no. 3 551–577.
- [8] G. T. Gilber, *Positive definite matrices and Sylvester’s criterion*, *Am. Math. Monthly* **98** (Jan., 1991) 44–46.
- [9] S. C. Frautschi and J. Kim, *SU(5) Higgs Problem With Adjoint + Vector Representations*, *Nucl.Phys.* **B196** (1982) 301.
- [10] J. Kim, *SU(N) Higgs Problem With Adjoint Representation and Michel’s Conjecture*, *Nucl.Phys.* **B197** (1982) 174.
- [11] J. Kim, *General Method for Analyzing Higgs Potentials*, *Nucl.Phys.* **B196** (1982) 285.
- [12] J. S. Kim, *SO(N) Higgs Problems with Adjoint + Vector Representations*, *Nucl.Phys.* **B207** (1982) 374.
- [13] J. S. Kim, *Orbit Spaces of Low Dimensional Representations of Simple Compact Connected Lie Groups and Extrema of a Group Invariant Scalar Potential*, *J.Math.Phys.* **25** (1984) 1694.
- [14] G. Sartori and G. Valente, *Allowed and observable phases in two Higgs doublet standard models*, [hep-ph/0304026](#).
- [15] M. Maniatis, A. von Manteuffel, O. Nachtmann, and F. Nagel, *Stability and symmetry breaking in the general two-Higgs-doublet model*, *Eur.Phys.J.* **C48** (2006) 805–823, [[hep-ph/0605184](#)].
- [16] I. Ivanov, *Minkowski space structure of the Higgs potential in 2HDM*, *Phys.Rev.* **D75** (2007) 035001, [[hep-ph/0609018](#)].

- [17] I. P. Ivanov and J. P. Silva, *Tree-level metastability bounds for the most general two Higgs doublet model*, *Phys. Rev.* **D92** (2015), no. 5 055017, [[arXiv:1507.05100](#)].
- [18] M. Maniatis and O. Nachtmann, *Stability and symmetry breaking in the general three-Higgs-doublet model*, *JHEP* **02** (2015) 058, [[arXiv:1408.6833](#)]. [Erratum: JHEP10,149(2015)].
- [19] M. Maniatis and O. Nachtmann, *Stability and symmetry breaking in the general n-Higgs-doublet model*, *Phys. Rev.* **D92** (2015), no. 7 075017, [[arXiv:1504.01736](#)].
- [20] A. Arhrib, R. Benbrik, M. Chabab, G. Moulhaka, M. C. Peyranere, L. Rahili, and J. Ramadan, *The Higgs Potential in the Type II Seesaw Model*, *Phys. Rev.* **D84** (2011) 095005, [[arXiv:1105.1925](#)].
- [21] C. Bonilla, R. M. Fonseca, and J. W. F. Valle, *Consistency of the triplet seesaw model revisited*, *Phys. Rev.* **D92** (2015), no. 7 075028, [[arXiv:1508.02323](#)].
- [22] G. Belanger, K. Kannike, A. Pukhov, and M. Raidal, *Impact of semi-annihilations on dark matter phenomenology - an example of Z_N symmetric scalar dark matter*, *JCAP* **1204** (2012) 010, [[arXiv:1202.2962](#)].
- [23] G. Bélanger, K. Kannike, A. Pukhov, and M. Raidal, *Minimal semi-annihilating Z_N scalar dark matter*, *JCAP* **1406** (2014) 021, [[arXiv:1403.4960](#)].
- [24] D. Eriksson, J. Rathsmann, and O. Stal, *2HDMC: Two-Higgs-Doublet Model Calculator Physics and Manual*, *Comput. Phys. Commun.* **181** (2010) 189–205, [[arXiv:0902.0851](#)].
- [25] K.P. and Hadeler, *On copositive matrices*, *Linear Algebra and its Applications* **49** (1983), no. 0 79 – 89.
- [26] K. Hadeler, *On copositive matrices*, *Linear Algebra and its Applications* **49** (1983), no. 0 79 – 89.
- [27] G. Chang and T. W. Sederberg, *Nonnegative quadratic Bézier triangular patches*, *Computer Aided Geometric Design* **11** (1994), no. 1 113 – 116.
- [28] R. Cottle, G. Habetler, and C. Lemke, *On classes of copositive matrices*, *Linear Algebra and its Applications* **3** (1970), no. 3 295 – 310.
- [29] W. Kaplan, *A test for copositive matrices*, *Linear Algebra and its Applications* **313** (July, 2000) 203–206.
- [30] M. Abud and G. Sartori, *The Geometry of Spontaneous Symmetry Breaking*, *Annals Phys.* **150** (1983) 307.
- [31] M. Abud and G. Sartori, *The Geometry of Orbit Space and Natural Minima of Higgs Potentials*, *Phys. Lett.* **B104** (1981) 147.
- [32] H. Weyl, *The Classical Groups, Their Invariants and Representation*. Princeton University Press, Princeton, NJ, 2nd ed., 1946.
- [33] J. A. Dieudonné and J. B. Carrell, *Invariant theory, old and new*, *Advances in Mathematics* **4** (1970), no. 1 1 – 80.
- [34] D. Luna, *Fonctions différentiables invariantes sous l’opération d’un groupe réductif*, *Annales de l’institut Fourier* **26** (1976), no. 1 33–49.
- [35] E. L. Rees, *Graphical discussion of the roots of a quartic equation*, *The American Mathematical Monthly* **29** (1922), no. 2 pp. 51–55.

- [36] D. Lazard, *Quantifier elimination: Optimal solution for two classical examples*, *Journal of Symbolic Computation* **5** (1988), no. 1–2 261 – 266.
- [37] G. Ulrich and L. T. Watson, *Positivity conditions for quartic polynomials*, tech. rep., Blacksburg, VA, USA, 1990.
- [38] P. Ko and Y. Tang, *Self-interacting scalar dark matter with local Z_3 symmetry*, *JCAP* **1405** (2014) 047, [[arXiv:1402.6449](#)].
- [39] I. F. Ginzburg and M. Krawczyk, *Symmetries of two Higgs doublet model and CP violation*, *Phys. Rev.* **D72** (2005) 115013, [[hep-ph/0408011](#)].
- [40] N. G. Deshpande and E. Ma, *Pattern of Symmetry Breaking with Two Higgs Doublets*, *Phys. Rev.* **D18** (1978) 2574.
- [41] K. Klimenko, *On Necessary and Sufficient Conditions for Some Higgs Potentials to Be Bounded From Below*, *Theor. Math. Phys.* **62** (1985) 58–65.
- [42] S. Nie and M. Sher, *Vacuum stability bounds in the two Higgs doublet model*, *Phys. Lett.* **B449** (1999) 89–92, [[hep-ph/9811234](#)].
- [43] S. Kanemura, T. Kasai, and Y. Okada, *Mass bounds of the lightest CP even Higgs boson in the two Higgs doublet model*, *Phys. Lett.* **B471** (1999) 182–190, [[hep-ph/9903289](#)].
- [44] L. Qi, *Eigenvalues of a real supersymmetric tensor*, *Journal of Symbolic Computation* **40** (2005), no. 6 1302 – 1324.
- [45] L.-H. Lim, *Singular Values and Eigenvalues of Tensors: A Variational Approach*, *ArXiv Mathematics e-prints* (July, 2006) [[math/0607648](#)].
- [46] L. Qi, *Symmetric nonnegative tensors and copositive tensors*, *Linear Algebra and its Applications* **439** (2013), no. 1 228 – 238, [[arXiv:1211.5642](#)].
- [47] Y. Song and L. Qi, *Necessary and sufficient conditions for copositive tensors*, *Linear and Multilinear Algebra* **63** (2015), no. 1 120–131, [[arXiv:1302.6084](#)].
- [48] Q. Ni, L. Qi, and F. Wang, *An eigenvalue method for testing positive definiteness of a multivariate form*, *Automatic Control, IEEE Transactions on* **53** (June, 2008) 1096–1107.